


Reduced Life Expectancy

Reduced Life Expectancy

Adverse Effects and Withdrawal Problems at Psychiatric Drugs:

Metabolic syndrome, cardiovascular diseases (especially in elderly people, children and adolescents), Takotsubo cardiomyopathy (especially at additional restraint), thrombosis, embolisms, stroke, diabetes, overweight, renal failure, neuroleptic malignant syndrome, malignant hyperthermia, agranulocytosis, asphyxia, tardive dyskinesia, hormone chaa-es combined with sexual disorders and neoplasm in the mammary glands, malformations in newborn, paradoxical or intrinsic suicidal



Reduced Life Expectancy

Measures to Reduce Psychiatric Patients' Mortality

Saint Camille Association, West African Republic of Benin:

Social support, including personal hygiene, food, clean clothes, work rehabilitation (learn to farm, sew, raise small animals, bake,

Measures to Reduce Psychiatric Patients' Mortality

La Cura and Associazione Penelope. Sicily, Italy

Measures to Reduce Psychiatric Patients' Mortality

Synthetics and Profit Orientation

“Assuming that our world is becoming ever more artificial and ‘man-made,’ and the demands by our modern performance-oriented society on our mental stability are constantly on the rise, wouldn’t it make sense to investigate every

(p. 17)-

.... Or Ecology and Humanistic Orientation

Until the final abolition of these drugs, a lot of people need help and support to withdraw from the drugs. Alternative systems and decentralized services to meet the needs of people experiencing mental health problems would minimize and in the long run make the use of synthetic and toxic psychiatric drugs needless" (p. 308).

Bach Jensen K. (2004). Detoxification—in the large and in the small: Towards a culture of respect. In: P. Lehmann (Ed.), *Coming off psychiatric drugs: Successful withdrawal from neuroleptics, antidepressants, lithium, carbamazepine and tranquilizers* (pp. 303-309). Berlin / Eugene, OR / Shrewsbury: Peter Lehmann Publishing (ebook 2013)

